

DUFACOC-200 PLUS W.S.P.

Composition:

Contains per g water-soluble powder:

Amprolium HCl..... 250 mg
Menadione sodium bisulphite..... 5 mg
Sulfaquinoxaline sodium 250 mg

Description:

Amprolium is an anti-coccidial drug with activity against *Eimeria tenella*, *E. necatrix*, and *E. acervulina*. Amprolium acts by interfering with thiamine metabolism in the parasite.

Sulfaquinoxaline is a chemotherapeutic with bacteriostatic action against many gram-negative and gram-positive bacteria. Its coccidiostatic spectrum of activity includes *Eimeria* spp. (*E. necatrix*, *E. maxima*, *E. acervulina*, *E. brunetti*, etc.).

Vitamin K in the form of menadione is added for the prevention of hemorrhagic disease. In combination with sulfaquinoxaline and amprolium, Vitamin K exhibits a synergistic action against most of the common species of the *Eimeria* parasite.

Indications:

Treatment and prevention of coccidiosis in poultry, lambs, calves and swine.

Contraindications:

Hypersensitivity to any of the active substances. Do not use in layers producing eggs for human consumption or animals with impaired kidney function.

Dosage and administration:

For oral administration in drinking water.

Poultry:

Therapeutic dose 100 g per 200 litres of water, for 5 – 7 days
(equivalent to 2500 chickens of 5 weeks of age or 1500 chickens of 10 weeks of age or 800 layers)

Prophylactic dose..... 100 g per 400 litres of water, for 3 – 4 days

Lambs, calves and swine 1 g per 10 kg body weight, for 5 days
preferably as a drench or in drinking water.

Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment.

Precautions and warnings:

Use caution when administering in drinking water during hot weather. Water consumption will increase, which may result in overdose.

Withdrawal times:

Meat and offal: 10 days

Storage conditions:

Store dark, dry and below 25°C. Do not freeze.

Packing:

100 g sachets and 1 kg jars.