

MULTIVIT EXTRA

Composition:

Contains per ml solution:

| | | | |
|--|-----------|---------------------------------------|---------|
| Retinyl palmitate (A) | 50.000 IU | Folic acid | 0.05 mg |
| Cholecalciferol (D3) | 12,000 IU | Ascorbic acid (C) | 20 mg |
| α -Tocopherol acetate (E) | 20 mg | Nicotinamide | 0.5 mg |
| Thiamine HCl (B1) | 1.5 mg | Dexpanthenol..... | 1 mg |
| Riboflavin sodium phosphate (B2) . | 0.1 mg | Menadione sodium bisulfite (K3) | 1 mg |
| Pyridoxine HCl (B6) | 0.25 mg | Biotin | 0.01 mg |
| Cyanocobalamin (B12) | 0.01 mg | | |

Indications:

Treatment and prevention of vitamin deficiencies in cattle, sheep, Swine and poultry, caused by:

- increased requirements due to stress or illness, growth, production or performance.
- decreased intake due to feed quality, inappetence or general weakness.

Dosage and administration:

For oral administration in drinking water.

Calves, bulls and lactating cows 20 ml per 100 litres of water

Chicks and broilers 12 ml per 1,000 litres of water

Layers 40 ml per 1,000 litres of water

Piglets 5 ml per 100 litres of water

Swine 10 ml per 100 litres of water

Lambs and sheep 5 ml per 100 litres of water

Shake well before use. Once diluted, Multivit Extra should be used immediately and refreshed every 24 hours for the duration of the treatment.

Withdrawal times:

None.

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

100 ml and 1 litre bottles.